# Central Coast

# GASTROENTEROLOGY

## Dr David N. Parkin

F.R.A.C.P. Prov. No. 0627397A Suite C5 Kanwal Medical Centre 654 Pacific Hwy, Kanwal 2259 P: 43935550 F: 0243141650

E: receptiondrparkin@gmail.com W: https://centralcoastgastro.com.au

## **BOWEL PREPARATION FOR COLONOSCOPY – PICOPREP ORANGE**

### **PREPARATION**

Please purchase **PICOPREP ORANGE** from your local pharmacy several days before your procedure.

#### TIMELINE

DAYS 3 & 2 BEFORE COLONOSCOPY	DAY BEFORE	DAY OF PROCEDURE
White food diet	Hospital will ring with admission time.	Clear fluids only till 2 hrs prior to admission.
(see pre-colonoscopy diet)		
	Follow bowel prep	You will be unable to drive
Do not eat any foods with skin, seeds, nuts, grains or high fibre	instructions including increased clear fluids	home. Please organise transport.

## **IMPORTANT:**

## **Anticoagulation medication**

Including Warfarin, Iscover, Plavix, Xarelto, Pradaxa and Eliquis should be withheld as instructed at the time of consultation.

Other medications (including Aspirin) should be taken as usual.

Diabetic Medications Tablets stop 2 days prior

Insulin Halve dose day before

Do not take on day of procedure bring with you to the hospital

# PRE-COLONOSCOPY DIET

#### YOU CAN EAT - WHITE FOODS

White bread (no whole seeds)

**English Muffins** 

Cornflakes, Rice Bubbles

Pork, Chicken, Fish

Peeled Fruit

**Peeled Vegetables** 

(no beetroot)

Potato, Mash Potato

**Hot Chips** 

Crackers (Saos or Jatz)

Eggs

Butter/Margarine

Cheese Custard Ice Cream

White Noodles

White Rice

### DO NOT EAT -FIBRE/GRAINS

Corn, peas, strawberries Skin on apple, pear, peach, nectarines (peeled fruit ok) Muesli, cereal, muesli bars Nuts or foods containing nuts

Jams/marmalades

Citrus fruit, grapes, tomatoes

Sundried tomatoes

Skin on vegetables (peeled

vegetables are ok)

Legumes e.g baked beans,

kidney beans, mustard

Sultanas, raisins

Brown rice Coconut

Sesame, sunflower, poppy

seeds

Casseroles containing lentils,

Barley, split peas Fruit yoghurt Fibre supplements

#### **CLEAR FLUIDS**

Water

Black tea or coffee (NO milk or

non dairy creamer)

Soft drinks

Sports drinks (Gatorade,

Powerade)

Cordial

Strained fruit juice without pulp, clear apple juice Yellow or orange jelly (NO blackcurrant, red, purple, green or blue drinks or jelly)

Clear ice

# **SAMPLE DIET MENU**

**BREAKFAST** 

OR

LUNCH

**EARLY DINNER** 

Cornflakes with milk

Chicken sandwich (white bread)

White fish, mashed potato

or rice

White toast and eggs

Vanilla ice cream

# **INSTRUCTIONS FOR PICOPREP ORANGE**

The day before your procedure

- LIGHT BREAKFAST ONLY COMPLETED BY 9AM (NO SKINS, SEEDS, NUTS OR GRAINS OR HIGH IN FIBRE)
- Then start clear fluids only

### DOSE 1: 2.00 PM (day BEFORE procedure)

Prepare Dose 1 (1 sachet) according to the leaflet in the PICOPREP ORANGE box. Chill if desired. Drink your first dose of PICOPREP ORANGE <u>SLOWLY</u> over 60 minutes (you may alternate with clear fluids). Follow with 500ml of clear fluids. Continue to drink clear fluids only.

### DOSE 2: 8.00PM (day BEFORE procedure)

Prepare Dose 2 (1 sachet) according to the leaflet in the PICOPREP ORANGE box. Drink your second dose of PICOPREP ORANGE\* <u>SLOWLY</u> over 60 minutes (you may alternate with clear fluids). Follow with 500 mL of clear fluids. Continue to drink clear fluids only.

#### DOSE 3: 5 hours BEFORE admission time

Prepare Dose 3 (1 sachet) according to the leaflet in the PICOPREP ORANGE box.

Drink your second dose of PICOPREP ORANGE <u>SLOWLY</u> over 60 minutes (you may alternate with clear fluids). Follow with 500 mL of clear fluids.

You should continue drinking a lot of varied clear fluids up until your **NIL BY MOUTH** time of **2 hrs before** admission.

#### RECOMMENDED CLEAR FLUIDS

 Water, black tea or coffee (NO milk or non-dairy creamer), soft drinks, sports drinks e.g. Gatorade, Powerade, cordial, strained fruit juice without pulp, clear apple juice, yellow or orange jelly (NO blackcurrant, red, purple, green or blue drinks or jelly), clear ice.

This product is designed to give you watery diarrhoea so stay close to a toilet. A protective cream or ointment (Lanolin, Vaseline) may be applied to your anal area before diarrhoea begins.

If you are unsure of any aspect of the preparation or are experiencing side effects please contact your Physician.

An Excellent Preparation



A Bad Preparation

